

DRUG FACTS

Active ingredient(s)

Purpose

Use(s)

- temporarily relieves minor aches and pains due to:
- minor pain of arthritis
- backache
- headache
- the common cold
- muscular aches
- menstrual cramps
- toothache
- temporarily reduces fever

Warnings

Do not use

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if you have

Ask a doctor before use if you have

- problems or serious side effects from taking pain relievers or fever reducers
- stomach problems that last or come back, such as heartburn, upset stomach, or stomach pain
- ulcers
- high blood pressure
- taken a diuretic
- reached age 60 or older
- bleeding problems
- heart or kidney disease

Ask a doctor or pharmacist before use if you are

Ask a doctor or pharmacist before use if you are

- taking any other drug containing an NSAID (prescription or nonprescription)
- taking a blood thinning (anticoagulant) or steroid drug
- under a doctor's care for any serious condition
- taking any other drug

When using this product

When using this product

- take with food or milk if stomach upset occurs
- long term continuous use may increase the risk of heart attack or stroke

Stop use and ask a doctor if

Stop use and ask a doctor if

- you feel faint, vomit blood, or have bloody or black stools. These are signs of stomach bleeding.
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing
- it feels like the pill is stuck in your throat
- you develop heartburn
- stomach pain or upset gets worse or lasts
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding

Keep out of reach of children

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
- the smallest effective dose should be used
- do not take longer than 10 days, unless directed by a doctor (see Warnings)
- drink a full glass of water with each dose

Adults and children 12 years and older	<ul style="list-style-type: none">take 1 tablet every 8 to 12 hours while symptoms lastfor the first dose you may take 2 tablets within the first hourdo not exceed 2 tablets in any 8- to 12-hour perioddo not exceed 3 tablets in a 24-hour period
Children under 12 years	<ul style="list-style-type: none">ask a doctor

Other information

- each tablet contains: sodium 20 mg
- store at 20-25°C (68-77°F). Avoid high humidity and excessive heat above 40°C (104°F).

PAIN RELIEVER / FEVER REDUCER (NSAID)

Drug Facts

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- asthma (wheezing)
- skin reddening
- facial swelling
- shock
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause stomach bleeding.

The chance is higher if you:

- are age 60 or older

- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing an NSAID (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**This product is not manufactured or distributed by Bayer HealthCare LLC, owner of the registered trademark Aleve[®].

Dist. by: **GOLDLINE LABORATORIES, INC.**

Miami, FL 33137

Dist. 1996 0908REV 21

Inactive ingredients

carnauba wax, FD&C blue no. 2 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, talc, titanium dioxide